

Poultris

Oral Suspension

Trimethoprim + Sulfamethoxazole



Definition:

It is an off white-yellow coloured oral suspension which includes Trimethoprim and Sulfamethoxazole.

Composition:

Each ml contains 80 mg Trimethoprim and 400 mg Sulfamethoxazole.

Indications:

Poultris is indicated in the treatment of infections caused by sensitive Gram (+) and Gram (-) bacteria on broiler, turkey, calf and also on lamb and kid which rumen activity haven't started yet.

For broiler and turkey; Coliseptisemia, salpingitis, cholera, coryza and staphylococ infections and other infections which are caused by the sensitive bacteria.

For calf, lamb and kids; Bacterial diarrhoea, gastro-enteritis, coliseptisemia, broncho-pneumonia, calf diphtheria, inflammation of umbilical cord, secondary bacterial infections caused by viral infections.

Dosage and administration:

Poultris is added in clean drinking water. Combine pharmacologic dosage of Poultris is 16-32 mg/ kg total weight/day. (trimethoprim 80 mg/ml + sulfametoxazole 400 mg/ml, total 480 mg/ml).

Dosage should be used 1-2 times a day for 3-5 days. Practically; for hen and turkey for which collective treatment is needed 15-30 ml/ 450 kg total weight/day for 3-5 days is sufficient.

For calf, lamb and kids which rumen activity haven't started yet, 0,5-1 ml/15 kg total weight/day for 3-5 days is sufficient. It is recommended to divide the daily dosage into two.

Packing:

200 ml glass bottles in cardboard box and 1 lt. plastic bottle without cardboard box.